

**WO0172706**

Publication Title:

SYNTHESIS

OF

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number  
**WO 01/72706 A1**

(51) International Patent Classification<sup>7</sup>: C07D 207/34,  
319/06, 405/06, A61K 31/40, A61P 3/06

Bangalore 561 229, Karnataka (IN). NAIR, Satheesh  
[IN/IN]; 20th K.M. Hosur Road, Hebbagodi, Bangalore  
561 229, Karnataka (IN).

(21) International Application Number: PCT/IN00/00030

(74) Agents: ANAND, Pravin et al.; Anand & Anand Advoca-  
tates, B-41, Nizamuddin East, New Delhi 110 013 (IN).

(22) International Filing Date: 28 March 2000 (28.03.2000)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): BIOCON  
INDIA LIMITED [IN/IN]; 20th K.M. Hosur Road, Heb-  
bagodi, Bangalore 561 229, Karnataka (IN).

(81) Designated States (national): AG, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,  
DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID,  
IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,  
LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL,  
PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,  
UA, UG, US, UZ, VN, YU, ZW.

(72) Inventors; and

(75) Inventors/Applicants (for US only): SAMBASIVAN,  
Ganesh [IN/IN]; 20th K.M. Hosur Road, Hebbagodi,  
Bangalore 561 229, Karnataka (IN). SRIDHARAN,  
Madhavan [IN/IN]; 20th K.M. Hosur Road, Hebbagodi,  
Bangalore 561 229, Karnataka (IN). PADUDEVAS-  
TANA, Sathyashanker [IN/IN]; 20th K.M. Hosur  
Road, Hebbagodi, Bangalore 561 229, Karnataka (IN).  
POORNAPRAJNA, Acharya [IN/IN]; 20th K.M. Hosur  
Road, Hebbagodi, Bangalore 561 229, Karnataka (IN).  
MATHEW, Joy [IN/IN]; 20th K.M. Hosur Road, Heb-  
bagodi, Bangalore 561 229, Karnataka (IN). SRINATH,  
Sumithra [IN/IN]; 20th K.M. Hosur Road, Hebbagodi,

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- with amended claims

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: SYNTHESIS OF [R-(R\*,R\*)]-2-(4-FLUOROPHENYL)-BETA.DELTA-DIHYDROXY-5-(1-METHYLETHYL)-3-PHENYL-4-[(PHENYLAMINO)CARBONYL]-1H-PYRROLE-1-HEPTANOIC ACID HEMI CALCIUM SALT (ATORVAS-TATIN)

(57) Abstract: The present invention discusses a novel process for the synthesis of [R-(R\*,R\*)]-2-(4-fluorophenyl)-B.D-dihy-  
droxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid hemi calcium salt by using 4-flu-  
oro- $\alpha$ -[2-methyl-1-oxopropyl]- $\gamma$ -oxo-N- $\beta$ -diphenylbenzene butanamide with (4R)-methyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-diox-  
ane-3-acetate. The compound so prepared is useful as inhibitors of the enzyme HMG-CoA reductase and are thus used as  
hypolipidemic and hypocholesterolemic agents.



WO 01/72706 A1

**TITLE OF THE INVENTION:**

SYNTHESIS OF [R-(R\*,R\*)]-2-(4-FLUOROPHENYL)-BETA,DELTA-DIHYDROXY-5-(1-METHYLETHYL)-3-PHENYL-4-[(PHENYLAMINO)CARBONYL]-1H-PYRROLE-1-HEPTANOIC ACID HEMI CALCIUM SALT (ATORVASTATIN)

5

**FILED OF THE INVENTION:**

This invention relates to a process for manufacturing R-(R\*,R\*)]-2-(4-fluorophenyl)-B,D-dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbonyl]-1H-pyrrole-1-heptanoic acid  
10 hemi calcium salt, atorvastatin and the novel intermediates produced during the course of manufacture. The said compound is useful as inhibitors of the enzyme HMG-CoA reductase and are thus useful as hypolipidemic and hypocholesterolemic agents.

**BACKGROUND OF THE INVENTION**

15 US Patent. No. 4,681,893, discloses a route using resolution of the racemic product using R (+)  $\alpha$ -methyl benzyl amine. US patent No. 5,003,080 discloses a synthetic route for the preparation of the chiral form of atorvastatin. The patent discloses a process for the preparation of the lactone or its salts by coupling an ester of (4R)-6-  
20 (2-aminoethyl)-2,2-dialkyl-1,3-dioxane-3-acetate with 4-fluoro- $\alpha$ -[2-methyl-1-oxopropyl] $\gamma$ -oxo-N- $\beta$ -diphenylbenzenebutaneamide followed by deprotection and hydrolysis to give the product. The product suffers from the fact ozonolysis is required as one of the steps for the synthesis of the amino ketal intermediate, which is hazardous  
25 for large scale preparation. The patent describes an alternate procedure wherein 4-fluoro- $\alpha$ -[2-methyl-1-oxopropyl] $\gamma$ -oxo-N- $\beta$ -diphenylbenzenebutaneamide is reacted with 3-amino propinaldehyde acetal followed by conventional procedures to give atorvastatin.

US patent No. 5,216,174, No. 5,097,045, No. 5,103,024, No. 5,124,482, No. 5,149,837, No. 5,155,251, No. 5,216,174, No. 5,245,047, No. 5,273,995, No. 5,248,793, and No. 5,397,792 describes various minor modifications in the procedure for the preparation of  
5 atorvastatin calcium salt.

Synthesis of esters of (4R)-6-(2-aminoethyl)-2,2-dialkyl-1,3-dioxane-3-acetate is an important part of the preparation of atorvastatin calcium. US patent 5,155,251 also discloses a synthetic route for the synthesis of (3R)-4-cyano-3-hydroxy butyric acid esters  
10 from (S)-3-hydroxy butyrolactone, which in turn is synthesized from a suitable carbohydrate substrate.

Other patents like 5,292,939, 5,319,110 and 5,374,773 discloses the preparation of 3,4-dihydroxybutyric acid. However, isolation of this highly water soluble compound or its lactone is not  
15 attempted.

Another multi step procedure starting from (S)-malic acid (J. org. Chem., 1981, 46, 4319) is reported. Esters of (S)-malic acid have also been used (Chem. Lett., 1984, 1389) for the synthesis of the hydroxy lactone involving  $\text{BMS-NaBH}_4$  reduction, followed by  
20 lactonization. While a six step procedure from D-isoascorbic acid is also reported (Syn., 1987, 570) but this process requires a silica gel chromatographic separation of the diastereomeric mixtures.

Optical resolution of the racemic hydroxylactones using lipase is disclosed in US patent 5,084,392 but this method suffers from poor  
25 enantiomeric excess and loss of the other active isomer.

Thus, these prior art procedures involves cumbersome reaction conditions or expensive starting materials, reagents which are difficult to handle or hazardous for scale up, coupled with a multi step procedure which results in poor overall yield.

The object of the present invention is to disclose an inexpensive, simple and scalable route for the synthesis of atorvastatin.

To achieve the said objective, this invention provides a process  
5 involves the synthesis of compound R-(R\*,R\*)]-2-(4-fluorophenyl)-  
B,D-dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)  
carbonyl]-1H-pyrrole-1-heptanoic acid hemi calcium salt (formula  
XXII) comprising:

- 10 a) reacting of compound of formula XVII with a compound  
of structure XIV in a non-polar solvent, to give an  
intermediate of structure XXI,
- b) deprotection, followed by hydrolysis of the compound of  
structure XXI followed by cyclization in ethyl acetate to  
give the lactone,
- 15 c) treating the lactone with  $\text{NH}_3$  to give the ammonium salt,  
and
- d) reacting the ammonium salt with  $\text{CaCl}_2$  gives the  
corresponding calcium salt of formula XXII.

20 The non-polar solvent in step (a) is selected from xylene or  
acetonitrile.

The de-protecting agent in step (b) is selected from moist silica  
or PTSA.

The ammonium salt in step (c) is prepared by treating with  
aqueous  $\text{NH}_3$  or methanolic  $\text{NH}_3$  or by passing  $\text{NH}_3$  gas to the  
25 compound taken in EtOAc, MeOH, isopropyl alcohol, diisopropyl  
ether or cyclohexane.

The compound of formula XVII is prepared by:

- a) reacting meldrum acid with isobutyryl chloride in the presence of base in  $\text{CH}_2\text{Cl}_2$  to give an intermediate of formula XVIII,
- b) reacting the intermediate of formula XVIII with base in  $\text{CH}_2\text{Cl}_2$  to give an intermediate of formula XIX,
- c) reacting the intermediate of formula XIX with benzaldehyde in the presence of base and toluene to give a compound of formula XX, and
- d) reacting of compound of formula XX with 4-fluorobenzaldehyde with a metal cyanide and polar solvent in the presence of a base to give the compound of formula XVII.

In step (b) the base is selected from aniline, pyridine, triethylamine, dimethylaniline, diethylisopropylamine.

- In step (c) the base is selected from piperidine, pyrrolidine or diisopropylethylamine.

In the metal cyanide in step (d) is selected from NaCN, CuCN, KCN, AgCN or tetraalkyl ammonium cyanide.

- The polar solvent in step (d) is selected from DMSO, DMF,  $\text{CH}_3\text{CN}$ .

The base in step (d) is selected from diisopropylethyl amine, 1-methylimidazole, pyridine.

The compound of (4R)-methyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-3-acetate (Formula XIV) used in step (a) is prepared by:

- a) reacting a compound of formula II with meldrum's acid followed by hydrolysis to give a compound of formula III,
- b) reacting a compound of formula III with thionyl chloride to give an acid chloride which is then reacted with a meldrum's acid followed by alcoholysis to give a compound of formula IV.

c) reducing a compound of formula IV to give a compound of formula V,

d) reducing a compound of formula V to give a dihydroxy ester of formula VI,

5 e) converting a compound of formula VI to give a compound of formula VII,

f) ammonolysis of a compound of formula VII to give a compound of formula VIII,

g) alternatively reacting the compound of formula VII with  
10 phthalimide salt to give a compound of formula XVI,

h) reacting a compound of formula XVI with hydrazine hydrate to give the compound of formula VIII, and

i) where  $R_5$  = *ter*-butyl in formula VIII, formula XIV is obtained

15  $R_4$  in step (a) is chosen from. OMs, OTs, Br, Cl or I.

The alcohol in step (b) is chosen from methanol, ethanol, isopropanol, benzyl alcohol.

The reducing agent in step (c) is baker's yeast or a reagent of formula  $R^*_2\text{BCl}$  where  $R^*_2$  is chosen from (-)  $\alpha$  pinene or (+)  $\alpha$ -pinene.

20 The reducing agent in step (d) is chosen from dimethoxyethylboron or triethoxyboron and sodium borohydride.

The protecting group for conversion in step (e) is chosen from acetone, benzophenone, acetophenone or benzaldehyde.

The reagent for ammonolysis in step (f) is chosen from aqueous  
25  $\text{NH}_3$ , methanolic  $\text{NH}_3$  or gaseous  $\text{NH}_3$  in a solvent chosen from EtOAc, cyclohexane, methanol, isopropanol or diisopropyl ether.

The phthalimide salt in step (g) is chosen from potassium, sodium or lithium.

The compound of (4R)-methyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-3-acetate (Formula XIV) used in step (a) is prepared by:

- a) reacting a compound of formula II with meldrum's acid followed by hydrolysis to give a compound of formula IX,
- 5 b) reacting a compound of formula IX with tert-butyl acetate with a base at -30 to -80°C to give a compound of formula X,
- c) reducing a compound of formula X to give a compound of formula XI,
- d) reducing a compound of formula XI to give a dihydroxy  
10 ester of formula XII,
- e) Converting a compound of formula XII to give a compound of formula XIII,
- f) ammonolysis of a compound of formula XIII to give a compound of formula XIV,
- 15 g) alternatively reacting the compound of formula XIII with phthalimide salt to give a compound of formula XV and
- h) reacting a compound of formula XV with hydrazine hydrate to give the compound of formula VIII

20  $R_4$  in step (a) is chosen from, OMs, OTs, Br, Cl or I.

The reducing agent in step (c) is baker's yeast or a reagent of formula  $R^*_2BCl$  where  $R^*_2$  is chosen from (-)  $\alpha$  pinene or (+)  $\alpha$ -pinene.

The reducing agent in step (d) is chosen from  
25 dimethoxyethylboron or triethoxyboron and sodium borohydride.

The protecting group for conversion in step (e) is chosen from acetone, benzophenone, acetophenone or benzaldehyde.



The reagent for ammonolysis in step (f) is chosen from aqueous  $\text{NH}_3$ , methanolic  $\text{NH}_3$  or gaseous  $\text{NH}_3$  in a solvent chosen from EtOAc, cyclohexane, methanol, isopropanol or diisopropyl ether.

The phthalimide salt in step (g) is chosen from potassium,  
5 sodium or lithium.

The intermediate compounds of formula XVIII, XIV, XV, VII & IX whenever prepared by the above process.

The intermediate of formula VII, wherein  $\text{R}_5$  is chosen from  $\text{C}_1$ - $\text{C}_5$  alkyl, benzyl, isopropyl, cyclohexyl or tert-butyl.

10 The intermediate of formula IX, wherein  $\text{R}_4$  is chosen from  $\text{CH}_2\text{Br}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{I}$ ,  $\text{CH}_2\text{OMs}$  or  $\text{CH}_2\text{Ots}$ .

#### DETAILED DESCRIPTION OF THE INVENTION

The process of the present invention in its first aspect is a new, improved, economical, and commercially feasible method for  
15 preparing HMG CoA reductase inhibitors of Formula XXII which are useful as inhibitors of the enzyme HMG CoA reductase and are thus useful as hypolipidemic or hypocholesterolemic agents is outlined in Scheme 5.

#### *Structure XXII*

20 The synthetic scheme for the synthesis of the amino ester of formula VIII is outlined in scheme 1

#### *Scheme - 1*

Thus, a halo acetic acid of formula I, where  $\text{X} = \text{Cl}$ ,  $\text{Br}$  or  $\text{I}$  is treated with sodium or potassium salt of trifluoromethane sulfonic  
25 acid or p-toluenesulfonic acid in the presence of base chosen from  $\text{Et}_3\text{N}$ , pyridine, N,N-dimethylaniline, N,N-diisopropylethyl amine, etc., in a solvent like,  $\text{CH}_2\text{Cl}_2$ ,  $\text{CH}_3\text{CN}$ , DMF etc., to give an intermediate which is then converted to an acid chloride by choosing a reagent from thionyl chloride, oxalyl chloride,  $\text{PCl}_5$  etc in the

presence of a base chosen from  $\text{Et}_3\text{N}$ , pyridine,  $\text{N,N}$ -dimethylaniline,  $\text{N,N}$ -diisopropylethyl amine, etc., in a solvent like,  $\text{CH}_2\text{Cl}_2$ ,  $\text{CH}_3\text{CN}$ , DMF etc., to give a compound of formula II, which is subsequently treated with meldrum's acid in presence of pyridine and  $\text{CH}_2\text{Cl}_2$  to give an acyl meldrum acid which is hydrolyzed to give a compound of formula III.

A  $\beta$ -keto acid of formula III is then converted to its acid chloride by reacting with a reagent chosen from  $\text{PCl}_5$ , oxalyl chloride, thionyl chloride etc., in the presence of a base chosen from pyridine,  $\text{Et}_3\text{N}$ , diisopropylethyl amine,  $\text{N,N}$ -dimethylaniline etc., in a solvent chosen from  $\text{CH}_2\text{Cl}_2$ ,  $\text{CH}_3\text{CN}$ , DMF etc. to give an acid chloride which is then reacted with meldrum acid as described earlier to give an acyl meldrum acid which on subsequent reaction with an alcohol  $\text{R}_5\text{OH}$

Where  $\text{R}_5 = \text{C}_1\text{-C}_6$  alkyl,  $\text{C}_3$  to  $\text{C}_6$ -cycloalkyl, benzyl etc., to give an intermediate of formula IV.

The diketo ester compound of formula IV is then reduced either with baker's yeast or with a reagent of formula  $\text{R}^*_2\text{BCl}$

Where  $\text{R}^*_2$  is chosen from  $(-)\text{-}\alpha\text{-pinene}$ ,  $(+)\text{-}\alpha\text{-pinene}$ , etc., to give a hydroxy keto ester of formula V. A hydroxy ketone ester of Formula V is treated with a borane reagent of formula  $\text{R}_2^{10}\text{BOCH}_3$  wherein  $\text{R}^{10}$  is a alkyl group of C1 to 3, for example, methoxydiethylborane in the absence of air and subsequent treatment with a metal hydride, such as, for example, sodium borohydride in a solvent, such as, for example, methanol, tetrahydrofuran, mixtures thereof, and the like at about  $0^\circ\text{C}$ . to about  $-110^\circ\text{C}$ . for about 5 hours followed by subsequent treatment with an acid, such as, for example, glacial acetic acid, and the like to afford a compound of Formula VI.

Preferably, the reaction is carried out with methoxydiethylborane under a nitrogen atmosphere and subsequent treatment with sodium borohydride in a mixture of methanol and tetrahydrofuran at about  $-20^{\circ}\text{C}$ . to about  $-78^{\circ}\text{C}$ . for about 5 hours followed by the addition of  
5 glacial acetic acid.

A 3,5-dihydroxy ester of Formula VI is treated with a ketal-forming reagent of  $\text{R}^1\text{COR}^2$  wherein  $\text{R}^1$  or  $\text{R}^2$  is independently from  $\text{C}_1$  to  $\text{C}_3$  alkyl, phenyl,  
10 benzyl, substituted phenyl etc., for example, a ketal-forming reagent selected from acetone, 2,2-dimethoxypropane, 2-methoxypropene, cyclopentanone, cyclohexanone, 1,1-dimethoxycyclopentane, 1,1-dimethoxycyclohexane, and the like or optionally an acetal forming reagent, for example, benzaldehyde, and the like in the presence of an  
15 acid, such as, for example, methanesulfonic acid, camphorsulfonic acid, paratoluenesulfonic acid, and the like, in the presence of excess reagent or in an inert solvent, such as, for example, dichloromethane, and the like at about  $0^{\circ}\text{C}$ . to about the reflux temperature of the reagent or solvent to afford a compound of Formula VI wherein  $\text{R}^1$   
20 and  $\text{R}^2$  are as defined above. Preferably, the reaction is carried out with a ketone forming reagent of Formula VII, for example, 2,2-dimethoxypropane and acetone in the presence of methanesulfonic acid at room temperature.

A compound of formula VII, which is then taken up in a  
25 solvent like, methanol, EtOAc, hexane or a mixture of these solvents and is converted to a compound of formula VIII by treating with gaseous  $\text{NH}_3$ , or by treating with aqueous  $\text{NH}_4\text{OH}$  solution or methanolic ammonia to give a compound of formula VIII


Optionally a compound of formula XIV can be prepared as described in scheme 2

*Scheme - 2*

5 Thus a compound of formula II where,  $R = CH_2Br, CH_2Cl, CH_2I, CH_2OMs, CH_2OTs, etc.,$  with meldrum acid in the presence of a base chosen from pyridine,  $Et_3N$ ,  $N,N$ -dimethylaniline,  $N,N$ -diisopropylethylamine etc in  $CH_2Cl_2$  to give an acyl meldrum acid of formula IX.

10 The acyl meldrum acid is taken in a solvent selected from hexane, THF, ether, or a mixture of hexane, THF or ether and was cooled to  $-30$  to  $-80^\circ C$ . A solution of tertiary butyl acetate and a base chosen from  $n-BuLi$ , LDA, Lithium pyrrolidide, etc., is added at a temperature between  $-30$  to  $-80^\circ C$  to afford the diketo ester compound of formula X.

The diketo ester compound of formula X is then reduced either with baker's yeast or with a reagent of formula



Where  $R^*_2$  is chosen from  $(-)\text{-}\alpha\text{-pinene}$ ,  $(+)\text{-}\alpha\text{-pinene}$ , etc., to give a  
20 hydroxy keto ester of formula XI. A hydroxy ketone ester of Formula XI is treated with a borane reagent of formula  $R_2^{10}BOCH_3$  wherein  $R^{10}$  is a alkyl group of C1 to 3, for example, methoxydiethylborane in the absence of air and subsequent treatment with a metal hydride, such as, for example, sodium borohydride in a  
25 solvent, such as, for example, methanol, tetrahydrofuran, mixtures thereof, and the like at about  $0^\circ C$ , to about  $-110^\circ C$ , for about 5 hours followed by subsequent treatment with an acid, like glacial acetic acid, and the like to afford a compound of Formula XII. Preferably, the reaction is carried out with methoxydiethylborane under a

nitrogen atmosphere and subsequent treatment with sodium borohydride in a mixture of methanol and tetrahydrofuran at about -20°C. to about -78°C. for about 5 hours followed by the addition of glacial acetic acid.

5 A 3,5-dihydroxy ester of Formula XII is treated with a ketal-forming reagent of



wherein  $R^1$  or  $R^2$  is independently from  $C_1$  to  $C_3$  alkyl, phenyl, benzyl, substituted phenyl etc., for example, a ketal-forming reagent  
10 selected from the group consisting of acetone, 2,2-dimethoxypropane, 2-methoxypropene, cyclopentanone, cyclohexanone, 1,1-dimethoxycyclopentane, 1,1-dimethoxycyclohexane, and the like or optionally an acetal forming reagent, for example, benzaldehyde, and the like in the presence of an acid, such as methanesulfonic acid,  
15 camphorsulfonic acid, paratoluenesulfonic acid, and the like, in the presence of excess reagent or in an inert solvent, such as, for example, dichloromethane, and the like at about 0°C. to about the reflux temperature of the reagent or solvent to afford a compound of Formula XIII wherein  $R^1$  and  $R^2$  are as defined above. Preferably, the  
20 reaction is carried out with a ketone forming reagent, for example, 2,2-dimethoxypropane and acetone in the presence of methanesulfonic acid at about room temperature.

A compound of formula XIII, which is then taken up in a solvent like, methanol, EtOAc, hexane or a mixture of these solvents  
25 and is converted to a compound of formula XIV by treating with gaseous  $NH_3$ , or by treating with aqueous  $NH_4OH$  solution or methanolic ammonia.

Alternatively, the intermediate of formula VII or XIII can be converted to VIII and XIV respectively as shown in scheme 3

*Scheme - 3*

where  $R_6$  is chosen from Li, K, Na, quaternary ammonium etc., to give the substituted phthalimide of formula XV or XVI which on treatment with hydrazine hydrate gives the desired amino compound.

- 5 A amino ester of Formula VIII or XIV is reacted with a diketone of Formula XVII wherein the process for the preparation of the compound of formula XVII is described in scheme 4

*Scheme - 4*

- 10 A compound of formula XVII is prepared as described in scheme 6, which comprises of reacting isobutyryl chloride and meldrum's acid in the presence of a base chosen from pyridine, triethylamine, diisopropylethyl amine, dimethylaniline etc in  $CH_2Cl_2$  at  $0-5^\circ C$  for about 18h to give an intermediate of formula XVIII.
- 15 Preferably the reaction is done in pyridine and  $CH_2Cl_2$  at  $0^\circ C$ . The acyl meldrum acid so obtained is then reacted with aniline in a solvent chosen from  $CH_2Cl_2$ , acetonitrile, toluene etc., at the reflux temperature of the solvent for about 12h to afford the amide of formula XIX. Preferably the reaction is carried out in  $CH_2Cl_2$  by
- 20 stirring at room temperature.

The keto amide is then reacted with benzaldehyde in the presence of a base chosen from aqueous NaOH in acetic acid, sodium acetate and acetic acid, pyrrolidine and acetic acid, piperidine and toluene, pyrrolidine and toluene etc., at the reflux temperature and by

25 removal of water for about 26h to give the methylenephenyl intermediate of formula XX.

The compound of formula XX is treated with 4-fluorobenzaldehyde in the presence of a catalyst chosen from metallic cyanide where the metal is Ag, K, Na, Cu, tetraalkylammonium etc..

or trimethylsilyl cyanide in a polar solvent chosen from DMSO, DMF, acetonitrile etc., at the reflux temperature of the solvent to give a compound of formula XVII. Preferably the reaction is carried out by reacting 4-fluorobenzaldehyde and sodium cyanide in DMSO at  
5 reflux temperature.

The diketone of formula XVII is reacted with the amino ester of formula VIII or XIV as described in Scheme 5 in the presence of a catalyst of Formula  $R_{12}CO_2H$  wherein  $R_{12}$  is chosen from trifluoromethane sulfonic acid, methane  
10 sulfonic acid, p-toluene sulfonic acid and a solvent or mixtures thereof such as, for example, acetonitrile, xylene, diisopropyl ether and the like for about 24 to about 48 hours from 5 to 10°C to about the reflux temperature of the solvent with the removal of water to afford a compound of Formula VII. Preferably, the reaction is carried out in  
15 the presence of methanesulfonic acid and a mixture of xylene-hexane at reflux for about 48 hours with the removal of water.

#### *Scheme – 5*

The compound of formula XXI is converted to atorvastatin  
20 calcium as shown in scheme 6

#### *Scheme – 6*

Which involves the deprotection of the ketal followed by hydrolysis of the ester to give the free acid which is converted to its  
25 ammonium salt by reacting with either  $NH_4OH$ , methanolic  $NH_3$  or by bubbling gaseous  $NH_3$  to the solution of carboxylic acid in a solvent chosen from a mixture of EtOAc, hexane, diisopropyl ether, isopropanol, cyclohexane and methanol. Preferably the intermediate of formula XXI is de-protected using moist silica in  $CH_2Cl_2$  at room

temperature over a period of 24h and is then hydrolyzed using methanolic sodium hydroxide and acidified using dil HCl to give the free acid which is converted to its ammonium salt by passing gaseous  $\text{NH}_3$  in EtOAc. The ammonium salt is then treated with calcium acetate to give atorvastatin calcium.

The invention will now be described with reference to the following example.

### Example 1

#### 1.1 Preparation of 4-methyl-3-oxo-N-phenylpetanamide (Formula XIX).

To a suspension of malonic acid (104g) in acetic anhydride (120mL) at room temperature, Conc.  $\text{H}_2\text{SO}_4$  (3mL) was added. The mixture was cooled to  $20^\circ\text{C}$  followed by the addition of acetone (80mL) drop wise. The contents were stirred at room temperature (15min) and kept at  $0-5^\circ\text{C}$  overnight and filtered. The solid was washed with cold water and cold acetone and dried. The crude material was recrystallized from acetone-hexane mixture.

Meldrum's acid (59g) was dissolved in  $\text{CH}_2\text{Cl}_2$  (200 mL) and cooled to  $0^\circ\text{C}$ . Pyridine (73mL) was added drop wise over a period of 30 min and the mixture was stirred for an additional 10 min. Isobutyryl chloride (44g) was added drop wise over a period of 30 min. and the mixture was stirred at  $0^\circ\text{C}$  for 1h followed by stirring at room temperature over night. The mixture was poured into 1.5N HCl containing crushed ice. The layers were separated and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (2x100mL). The combined extracts were washed with 1.5N HCl (2x100mL) followed by saturated  $\text{NH}_4\text{Cl}$  solution (2x100mL) and dried over  $\text{Na}_2\text{SO}_4$  and concentrated under reduced pressure to afford the crude acyl meldrum's acid which was used for the next step.



The crude acyl meldrum's acid (84g) was taken in benzene (300mL) and aniline (111mL) was added. The mixture was refluxed for 4h. Cool the reaction mixture to room temperature and wash with 2N HCl (5x100mL) and benzene was removed under reduced  
5 pressure to get formula XIX.

### **1.2 Preparation of 4-methyl-3-oxo-N-phenyl-2-(phenylmethylene) pentanamide (Formula XX).**

The crude amide was taken in acetic acid and piperidine. To this  
10 mixture at room temperature benzaldehyde was added. The contents were allowed to stir under reflux for 2h. Pour the contents into water and extract with CH<sub>2</sub>Cl<sub>2</sub>. The organic extracts were washed with bicarbonate, bisulfite solution, dried and concentrated under reduced pressure to afford the crude compound of formula XX.

15

### **1.3 Preparation of 4-fluoro- $\alpha$ -[2-methyl-1-oxopropyl]- $\gamma$ -oxo-N- $\beta$ -diphenylbenzenebutaneamide (Formula XVII).**

To 4-fluorobenzaldehyde in anhydrous DMF, sodium cyanide was added and the contents were refluxed for 4h. To this the  
20 intermediate from example 3 was added and the contents were stirred for an additional 18h. Usual work up affords the crude diketo compound of formula XVII.

### **1.4 Preparation of 4-trifluoromethanesulfonyl-3-oxo-butyric acid (Formula III).**

25 To a solution of chloroacetic acid in pyridine at 0°C and CH<sub>2</sub>Cl<sub>2</sub>, trifluoromethane sulfonyl chloride was added slowly. After the reaction was complete, oxalyl chloride was added and the contents

were refluxed for 3h and the solvent was removed under reduced pressure to give the crude acid chloride.

Meldrum's acid (59g) was dissolved in  $\text{CH}_2\text{Cl}_2$  (200 mL) and cooled to  $0^\circ\text{C}$ . Pyridine (73mL) was added drop wise over a period of 30 min and the mixture was stirred for an additional 10 min. Trifluoromethanesulfonyl acetylchloride (44g) was added drop wise over a period of 30 min. and the mixture was stirred at  $0^\circ\text{C}$  for 1h followed by stirring at room temperature over night. The mixture was poured into 1.5N HCl containing crushed ice. The layers were separated and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (2x100mL). The combined extracts were washed with 1.5N HCl (2x100mL) followed by saturated  $\text{NH}_4\text{Cl}$  solution (2x100mL) and dried over  $\text{Na}_2\text{SO}_4$  and concentrated under reduced pressure to afford the crude acyl meldrum's acid was hydrolyzed to give the title compound of formula III.

#### 1.5 Preparation of methyl 6-trifluoromethane sulfonyl-5,3-dioxo-butryate (Formula IV).

The crude compound from example 5 was taken in  $\text{CH}_2\text{Cl}_2$  and pyridine and was cooled to  $0-5^\circ\text{C}$ . Oxalyl chloride was slowly introduced and the contents were refluxed for 6h. The solvent was removed under reduced pressure to afford the crude acid chloride.

Meldrum's acid (59g) was dissolved in  $\text{CH}_2\text{Cl}_2$  (200 mL) and cooled to  $0^\circ\text{C}$ . Pyridine (73mL) was added drop wise over a period of 30 min and the mixture was stirred for an additional 10 min. The acid chloride (44g) was added drop wise over a period of 30 min. and the mixture was stirred at  $0^\circ\text{C}$  for 1h followed by stirring at room temperature over night. The mixture was poured into 1.5N HCl containing crushed ice. The layers were separated and the aqueous

layer was extracted with  $\text{CH}_2\text{Cl}_2$  (2x100mL). The combined extracts were washed with 1.5N HCl (2x100mL) followed by saturated  $\text{NH}_4\text{Cl}$  solution (2x100mL) and dried over  $\text{Na}_2\text{SO}_4$  and concentrated under reduced pressure to afford the crude acyl meldrum's acid was  
5 refluxed in methanol to give the title compound of formula IV

**1.6 reparation of methyl 6-trifluoromethane sulfonyl-5-oxo-3-hydroxybutryate (Formula V).**

To a mixture of dry bakers' yeast and glucose in boiled water  
10 was stirred at ambient temperature. The crude diketo ester from example 5 was then added and the reaction mixture was stirred for an additional period 8-13h at ambient temperature. Celite was added to the reaction mixture and filtered. The celite bed was washed thoroughly with ethyl acetate and the combined ethyl acetate extracts  
15 was dried and concentrated under reduced pressure to afford the title product which was used for the next step without further purification.

**1.7 Preparation of (4R)-methyl 6-trifluoromethyl sulfonyl methyl-2,2-dimethyl-1,3-dioxane-4-acetate (Formula VII).**

20 The crude product from example 6 was taken up in dry THF and isopropanol under nitrogen atmosphere. The solution was cooled to  $-78^\circ\text{C}$  and a solution of diethylmethoxyborane was added. The reaction mixture was cooled to  $-80^\circ\text{C}$  and sodium borohydride was added in portion over a period of 4h. The temperature was maintained  
25 between  $-70$  to  $-85^\circ\text{C}$  and was allowed to warm to room temperature and stand for 18h. The reaction was quenched by addition of acetic acid and concentrated under reduced pressure to afford an oily residue.

The crude oil was taken up in acetone and 2,2-dimethoxypropane was added followed by catalytic amount of trifluoromethanesulfonic acid. The contents were stirred at room temperature for 4h and after the completion of the reaction was washed with bicarbonate solution. The organic extracts was washed with water and was concentrated to give an oil which was crystallized from hexane.

**1.8 Preparation of (4R)-methyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-3-acetate (Formula VIII)**

The crude mesylate from example 8 was taken up in methanol and  $\text{NH}_3$  gas was bubbled. The contents were then stirred for 6-14h at room temperature. After the reaction was complete, solvent was removed under reduced pressure to afford the crude title compound of formula VIII.










**1.9 Preparation of [R-(R\*,R\*)]-2-(4-fluorophenyl)- $\beta$ , $\delta$ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[phenylaminocarbonyl]-1H-pyrrole-1-heptanoic acid, hemi calcium salt (Formula XXII)**

A solution of (4R)-methyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-3-acetate (of formula XIV) and 4-fluoro- $\alpha$ -[2-methyl-1-oxopropyl] $\gamma$ -oxo-N- $\beta$ -diphenylbenzenebutaneamide (formula XVII) and acetic acid in xylene were heated to reflux to 44h. The solution was diluted with diisopropyl ether and methanol and was washed with dilute methanolic sodium hydroxide solution, dilute HCl and the solvent was then removed under vacuum. The crude oil was stirred with moist silica in  $\text{CH}_2\text{Cl}_2$  and was stirred at room temperature for 18h. A solution of aqueous NaOH was then added at room temperature and was stirred for 4h. The reaction mixture was diluted



with water and was washed with diisopropyl ether. The aqueous layer was acidified with HCl and was taken up in diisopropyl ether. The crude acid intermediate was then taken up in EtOAc and  $\text{NH}_3$  gas was bubbled. The contents were stirred for completion of the reaction and  
5 solvent was removed upon which the product crystallized. The crude ammonium salt is then taken up in diisopropyl ether-isopropanol mixture and a solution of calcium acetate was added at room temperature upon which the calcium salt precipitated from the solution. The product was filtered and dried under vacuum to get  
10 formula XXII of acceptable pharmaceutical purity.









The invention has been described by reference to specific embodiments, this was for the purpose of illustration only. Numerous alternative embodiments will be apparent to those skilled in the art and are considered within the scope of these claims.



20




10








5






### Scheme -1




5

### Scheme - 2




Scheme - 3




Scheme - 4

5




25

Scheme - 5



5

Scheme - 6



10

**We claim:**

1. The process involves the synthesis of compound R-(R\*,R\*)]-2-(4-fluorophenyl)-B,D-dihydroxy-5-(1-methylethyl)-3-phenyl-4-  
5 [(phenylamino) carbonyl]-1H-pyrrole-1-heptanoic acid hemi calcium salt (formula XXII) which comprising:
  - a) reacting of compound of formula XVII with a compound of structure XIV in a non-polar solvent, to give an intermediate of structure XXI,
  - 10 b) deprotection, followed by hydrolysis of the compound of structure XXI followed by cyclization in ethyl acetate to give the lactone,
  - c) treating the lactone with NH<sub>3</sub> to give the ammonium salt, and
  - 15 d) reacting the ammonium salt with CaCl<sub>2</sub> gives the corresponding calcium salt of formula XXII.
2. A process as claimed in claim 1, wherein the non-polar solvent in step (a) is selected from xylene or acetonitrile.
3. A process as claimed in claim 1, wherein the de-protecting  
20 agent in step (b) is selected from moist silica or PTSA.
4. A process as claimed in claim 1, wherein the ammonium salt in step (c) is prepared by treating with aqueous NH<sub>3</sub> or methanolic NH<sub>3</sub> or by passing NH<sub>3</sub> gas to the compound taken in EtOAc, MeOH, isopropyl alcohol, diisopropyl ether or cyclohexane.
- 25 5. A process as claimed in claim 1, wherein the compound of compound of formula XVII is prepared by:
  - a) reacting meldrum acid with isobutyryl chloride in the presence of base in CH<sub>2</sub>Cl<sub>2</sub> to give an intermediate of formula XVIII,

b) reacting the intermediate of formula XVIII with base in  $\text{CH}_2\text{Cl}_2$  to give an intermediate of formula XIX,

c) reacting the intermediate of formula XIX with benzaldehyde in the presence of base and toluene to give a compound  
5 of formula XX, and

d) reacting of compound of formula XX with 4-fluorobenzaldehyde with a metal cyanide and polar solvent in the presence of a base to give the compound of formula XVII.

6. A process as claimed in claim 5 where in step (b) the base is  
10 selected from aniline, pyridine, triethylamine, dimethylaniline, diethylisopropylamine.

7. A process as claimed in claim 5 where in step (c) the base is selected from piperidine, pyrrolidine or diisopropylethylamine.

8. A process as claimed in claim 5 where in the metal cyanide in  
15 step (d) is selected from NaCN, CuCN, KCN, AgCN or tetraalkyl ammonium cyanide.

9. A process as claimed in claim 5 wherein the polar solvent in step (d) is selected from DMSO, DMF,  $\text{CH}_3\text{CN}$ .

10. A process as claimed in claim 5 wherein the base in step (d) is  
20 selected from diisopropylethyl amine, 1-methylimidazole, pyridine.

11. A process as claimed in claim 1 wherein compound of (4R)-methyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-3-acetate (Formula XIV) used in step (a) is prepared by:

a) reacting a compound of formula II with meldrum's acid  
25 followed by hydrolysis to give a compound of formula III,

b) reacting a compound of formula III with thionyl chloride to give an acid chloride which is then reacted with a meldrum's acid followed by alcoholysis to give a compound of formula IV,

- c) reducing a compound of formula IV to give a compound of formula V,
  - d) reducing a compound of formula V to give a dihydroxy ester of formula VI,
  - 5 e) Converting a compound of formula VI to give a compound of formula VII,
  - f) ammonolysis of a compound of formula VII to give a compound of formula VIII,
  - g) alternatively reacting the compound of formula VII with  
10 phthalimide salt to give a compound of formula XVI,
  - h) reacting a compound of formula XVI with hydrazine hydrate to give the compound of formula VIII, and
  - i) where  $R_5$  = ter-butyl in formula VIII, formula XIV is obtained
- 15 12. A process as claimed in claim 11 wherein  $R_4$  in step (a) is chosen from, OMs, OTs, Br, Cl or I.
13. A process as claimed in claim 11 wherein, the alcohol in step (b) is chosen from methanol, ethanol, isopropanol, benzyl alcohol.
14. A process as claimed in claim 11 wherein the reducing agent in  
20 step (c) is baker's yeast or a reagent of formula  $R^*_2BCl$  where  $R^*_2$  is chosen from (-)  $\alpha$  pinene or (+)  $-\alpha$ - pinene.
15. A process as claimed in claim 11 wherein the reducing agent in step (d) is chosen from dimethoxyethylboron or triethoxyboron and sodium borohydride.
- 25 16. A process as claimed in claim 11 wherein the protecting group for conversion in step (e) is chosen from acetone, benzophenone, acetophenone or benzaldehyde.
17. A process as claimed in claim 11 wherein the reagent for ammonolysis in step (f) is chosen from aqueous  $NH_3$ , methanolic  $NH_3$

or gaseous  $\text{NH}_3$  in a solvent chosen from EtOAc, cyclohexane, methanol, isopropanol or diisopropyl ether.

18. A process as claimed in claim 11 wherein the phthalimide salt in step (g) is chosen from potassium, sodium or lithium.

5 19. A process as claimed in claim 1 wherein compound of (4R)-methyl 6-(2-aminoethyl)-2,2-dimethyl-1,3-dioxane-3-acetate (Formula XIV) used in step (a) is prepared by:

a) reacting a compound of formula II with meldrum's acid followed by hydrolysis to give a compound of formula IX,

10 b) reacting a compound of formula IX with tert-butyl acetate with a base at  $-30$  to  $-80^\circ\text{C}$  to give a compound of formula X,

c) reducing a compound of formula X to give a compound of formula XI,

15 d) reducing a compound of formula XI to give a dihydroxy ester of formula XII,

e) Converting a compound of formula XII to give a compound of formula XIII,

f) ammonolysis of a compound of formula XIII to give a compound of formula XIV

20 g) alternatively reacting the compound of formula XIII with phthalimide salt to give a compound of formula XV and

h) reacting a compound of formula XV with hydrazine hydrate to give the compound of formula XIV.

25 20. A process as claimed in claim 19 wherein  $\text{R}_4$  in step (a) is chosen from, OM, OT, Br, Cl or I.

21. A process as claimed in claim 19 wherein the reducing agent in step (c) is baker's yeast or a reagent of formula  $\text{R}^*_2\text{BCl}$  where  $\text{R}^*_2$  is chosen from (-)  $\alpha$  pinene or (+)  $\alpha$ - pinene.

22. A process as claimed in claim 19 wherein the reducing agent in step (d) is chosen from dimethoxyethylboron or triethoxyboron and sodium borohydride.
23. A process as claimed in claim 19 wherein the protecting group  
5 for conversion in step (e) is chosen from acetone, benzophenone, acetophenone or benzaldehyde.
24. A process as claimed in claim 19 wherein the reagent for ammonolysis in step (f) is chosen from aqueous  $\text{NH}_3$ , methanolic  $\text{NH}_3$  or gaseous  $\text{NH}_3$  in a solvent chosen from EtOAc, cyclohexane,  
10 methanol, isopropanol or diisopropyl ether.
25. A process as claimed in claim 19 wherein the phthalimide salt in step (g) is chosen from potassium, sodium or lithium.
26. The intermediate compound of formula XVIII as claimed in claim 5.
- 15 27. The intermediate of formula XIV as claimed in claim 1.
28. The intermediate of formula XV as claimed in claim 19.
29. The intermediate of formula VII, wherein  $\text{R}_5$  is chosen from  $\text{C}_1\text{-C}_5$  alkyl, benzyl, isopropyl, cyclohexyl or tert-butyl as claimed in claim 11.
- 20 30. The intermediate of formula IX, wherein  $\text{R}_4$  is chosen from  $\text{CH}_2\text{Br}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{I}$ ,  $\text{CH}_2\text{OMs}$  or  $\text{CH}_2\text{Ots}$  as claimed in claim 19.



## AMENDED CLAIMS

[received by the International Bureau on 16 February 2001 (16.02.01);  
original claims 1-30 replaced by new claims 1-21 (5 pages)]

- 5 1. The process for the manufacture of compound R-(R\*, R\*)]-2-) 4-fluorophenyl)-  
β,δ-dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbonyl]-1H-  
pyrrole-1-heptanoic acid hemi calcium salt (formula XXII) which comprises:
- 10 a) reacting of compound of formula XVII with a compound of structure  
XIV or a compound of structure VIII in a mixture of solvents chosen  
from xylene, cyclohexane, methyl tert-butyl ether, diisopropyl ether,  
acetonitrile, in the presence of catalyst chosen from trifluoromethyl  
15 sulfonic acid, methane sulfonic acid, p-toluene sulfonic acid (PTSA) or  
pyridine p-toluene sulphonic acid (PPTS), to give an intermediate of  
structure XXI, where R<sub>1</sub> or R<sub>2</sub> is independently from H, C<sub>1</sub> to C<sub>3</sub> alkyl,  
phenyl, benzyl, substituted phenyl and R<sub>3</sub> is chosen from C<sub>1</sub>-C<sub>6</sub> n-alkyl,  
C<sub>3</sub>-C<sub>6</sub> Cycloalkyl, benzyl or ter-butyl,
- 20 b) deprotection followed by hydrolysis of the compound of structure XXI  
followed by cyclization in ethyl acetate to give the lactone,
- c) treating the lactone with NH<sub>3</sub> to give the ammonium salt, and
- 25 d) reacting the ammonium salt with Ca<sup>+2</sup> salts gives the corresponding  
calcium salt of formula XXII.
2. A process as claimed in claim 1, wherein the ammonium salt in step (c) is  
30 prepared by treating with aqueous NH<sub>3</sub> or methanolic NH<sub>3</sub> or by passing NH<sub>3</sub>  
gas to the compound taken in EtOAc, MeOH, Isopropyl alcohol, diisopropyl  
ether or cyclohexane.
3. A process as claimed in claim 1, wherein the compound of formula XVII is  
35 prepared by:
- a) reacting meldrum acid with isobutyryl chloride in the presence of base in  
CH<sub>2</sub>Cl<sub>2</sub> to give an intermediate of formula XVIII.
- 40 b) reacting the intermediate of formula XVIII with aniline in CH<sub>2</sub>Cl<sub>2</sub> to give  
an intermediate of formula XIX,

5 c) reacting the intermediate of formula XIX with benzaldehyde in the presence of catalyst and toluene to give a compound of formula XX, and

d) reacting of compound of formula XX with 4-fluorobenzaldehyde with a cyanide reagent and solvent in the presence of a base to give the compound of formula XVII.

10 4. A process as claimed in claim 3 wherein step (c) the catalyst is selected from piperidine, pyrrolidine, diisopropylethylamine, PTSA or PPTS.

15 5. A process as claimed in claim 3 where in the cyanide reagent in step (d) is selected from NaCN, CuCN, KCN, AgCN, tetra-alkyl ammonium cyanide or trimethyl silyl cyanide.

6. A process as claimed in claim 3 wherein the solvent in step (d) is selected from DMSO, DMF, CH<sub>3</sub>CN.

20 7. A process as claimed in claim 3 wherein the base in step (d) is selected from diisopropylethyl amine, 1-methylimidazole, pyridine.

25 8. A process as claimed in claim 1 wherein compound of formula XIV used in step (a); where R<sub>1</sub> or R<sub>2</sub> is independently from H, C<sub>1</sub> to C<sub>3</sub> alkyl, phenyl, benzyl, substituted phenyl and R<sub>4</sub> is chosen from CH<sub>2</sub>OMs, CH<sub>2</sub>OTs, CH<sub>2</sub>Cl, CH<sub>2</sub>Br or CH<sub>2</sub>I; is prepared by:

30 a) reacting a compound of formula II, with meldrum's acid followed by hydrolysis to give a compound of formula IX,

b) reacting a compound of formula IX with ter-butyl acetate with a base at -30 to -80 °C to give a compound of formula X,

35 c) reducing a compound of formula X to give a compound of formula XI.

d) reducing a compound of formula XI to give a dihydroxy ester of formula XII,

40 e) converting a compound of formula XII to give a compound of formula XIII.

- 5 f) ammonolysis of a compound of formula XIII to give a compound of formula XIV.
- g) alternatively reacting the compound of formula XIII with phthalimide salt to give a compound of formula XV, and
- 10 h) reacting a compound of formula XV with hydrazine hydrate to give the compound of formula XIV.
9. A process as claimed in claim 8 wherein the reducing agent in step (c) is derived from yeast or a reagent of formula  $R^*_2BCl$ , where  $R^*_2$  is chosen from  $\alpha$  pinene.
- 15 10. A process as claimed in claim 8 wherein the reducing agent in step (d) is chosen from dimethoxyethylboron or triethoxyboron and sodium borohydride.
- 20 11. A process as claimed in claim 8 wherein the reagent for ammonolysis in step (f) is chosen from aqueous  $NH_3$ , methanolic  $NH_3$  or gaseous  $NH_3$  in a solvent chosen from EtOAc, cyclohexane, methanol, isopropanol or diisopropyl ether.
- 25 12. A process as claimed in claim 8 wherein the phthalimide salt in step (g) is chosen from potassium, sodium or lithium.
- 30 13. A process as claimed in claim 1 wherein compound of formula VIII used in step (a); where  $R_1$  or  $R_2$  is independently from H,  $C_1$  to  $C_3$ , alkyl, phenyl, benzyl, substituted phenyl  $R_4$  is chosen from  $CH_2OMs$ ,  $CH_2OTs$ ,  $CH_2Cl$ ,  $CH_2Br$  or  $CH_2I$  and  $R_5$  is  $C_1$ - $C_6$  n-alkyl,  $C_3$ - $C_6$  cycloalkyl, benzyl or ter-butyl; is prepared by:
- 35 a) reacting a compound of formula II, with meldrum's acid followed by hydrolysis to give a compound of formula III,
- b) reacting a compound of formula III with thionyl chloride to give an acid chloride which is then reacted with a meldrum's acid followed by alcoholysis to give a compound of formula IV,
- 40 c) reducing a compound of formula IV to give a compound of formula V.

- 5 d) reducing a compound of formula V to give a dihydroxy ester of formula VI,
- e) converting a compound of formula VI to give a compound of formula VII.
- 10 f) ammonolysis of a compound of formula VII to give a compound of formula VIII.
- g) alternatively reacting the compound of formula VII with phthalimide salt to give a compound of formula XVI, and
- 15 h) reacting a compound of formula XVI with hydrazine hydrate to give the compound of formula VIII.

20 14. A process as claimed in claim 13 wherein the reducing agent in step (c) is derived from yeast or a reagent of formula  $R^*_2BCl$ , where  $R^*_2$  is chosen from  $\alpha$  pinene.

25 15. A process as claimed in claim 13 wherein the reducing agent in step (d) is chosen from dimethoxyethylboron or tetroxyboron and sodium borohydride.

16. A process as claimed in claim 13 wherein the reagent for ammonolysis in step (f) is chosen from aqueous  $NH_3$ , methanolic  $NH_3$  or gaseous  $NH_3$  in a solvent chosen from EtOAc, cyclohexane, methanol, isopropanol or diisopropyl ether.

30 17. A process as claimed in claim 13 wherein the phthalimide salt in step (g) is chosen from potassium, sodium or lithium.

35 18. The intermediate of formula XVI, where  $R_1$  or  $R_2$  is independently from H,  $C_1$  to  $C_3$  alkyl, phenyl, benzyl, substituted phenyl and  $R_5$  is chosen from  $C_1 - C_6$  n-alkyl,  $C_3 - C_6$  cycloalkyl, benzyl or ter-butyl.

19. The intermediate of formula XV, where  $R_1$  or  $R_2$  is independently from H,  $C_1$  to  $C_3$  alkyl, phenyl, benzyl, substituted phenyl.

20. The intermediate of formula VII, where  $R_1$  or  $R_2$  is independently from H,  $C_1$  to  $C_3$  alkyl, phenyl, benzyl, substituted phenyl and  $R_4$  is chosen from  $CH_2Br$ ,  $CH_2Cl$ ,  $CH_2I$ ,  $CH_2OMs$  or  $CH_2OTs$  and  $R_5$  is chosen from  $C_1 - C_6$  n alkyl,  $C_3 - C_6$  cycloalkyl, benzyl or ter-butyl.
21. The intermediate of formula IX, wherein  $R_4$  is chosen from  $CH_2Br$ ,  $CH_2Cl$ ,  $CH_2I$ ,  $CH_2OMs$  or  $CH_2OTs$ .

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/IN 00/00030

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D207/34 C07D319/06 C07D405/06 A61K31/40 A61P3/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BAUMANN K L ET AL: "THE CONVERGENT SYNTHESIS OF CI-981, AN OPTICALLY ACTIVE, HIGHLY POTENT, TISSUE SELECTIVE INHIBITOR OF HMG-COA REDUCTASE" TETRAHEDRON LETTERS,NL,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, vol. 33, no. 17, 21 April 1992 (1992-04-21), pages 2283-2284, XP000608147 ISSN: 0040-4039 the whole document	1-10
X	EP 0 330 172 A (WARNER LAMBERT CO) 30 August 1989 (1989-08-30) cited in the application Scheme III and V page 6, line 44 -page 11, line 18 --- -/--	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

15 November 2000

Date of mailing of the international search report

24/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Seitner, I

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IN 00/00030

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ZAWACKI F J ET AL: "A Convenient Synthesis of Unsymmetrical, Substituted gamma-Pyrones from Meldrum's Acid" TETRAHEDRON LETTERS,NL,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, vol. 37, no. 36, 2 September 1996 (1996-09-02), pages 6499-6502, XP004030727  ISSN: 0040-4039  Scheme 2 compound 5  -----</p>	26

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/IN 00 00030

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 11-18, 19-25, 27-29

Claims 11-18, 19-25, and 27-29 do not meet the requirements of Article 6 PCT in that the matter for which protection is sought is not clearly defined. The following functional statements do not enable the skilled person to determine which technical features are necessary to perform the stated functions:

In claims 11-18 no definitions have been provided for R1, R2, R4, and/or R5.

In claims 19-25 no definitions have been provided for R1, R2, and/or R4.

In claims 12 and 20 the definitions of R4 seem to be incorrect.

In claims 27-29 R1 and R2 have not been defined.

Furthermore, in claim 29 R4 has not been defined.

Due to the lack of clarity within the meaning of Article 6 PCT a meaningful search of the claims 11-18, 19-25, 27-29 is impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear, namely claims 1-10 and claims 26 and 30.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IN 00/00030

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0330172 A	30-08-1989	US 5003080 A	26-03-1991
		AT 109777 T	15-08-1994
		AU 634689 B	25-02-1993
		AU 1601792 A	09-07-1992
		AU 635171 B	11-03-1993
		AU 1601892 A	09-07-1992
		AU 3349689 A	06-09-1989
		AU 621874 B	26-03-1992
		CA 1330441 A	28-06-1994
		DE 68917336 D	15-09-1994
		DE 68917336 T	01-12-1994
		DK 197090 A	04-10-1990
		EP 0448552 A	02-10-1991
		ES 2058356 T	01-11-1994
		FI 94958 B	15-08-1995
		FI 941550 A,B,	05-04-1994
		HK 1000732 A	24-04-1998
		IE 63994 B	28-06-1995
		JP 3009139 B	14-02-2000
		JP 10195071 A	28-07-1998
		JP 2843627 B	06-01-1999
		JP 3502798 T	27-06-1991
		KR 9711578 B	12-07-1997
		KR 9711579 B	12-07-1997
		KR 123813 B	27-11-1997
		KR 9711462 B	11-07-1997
		KR 137884 B	01-05-1998
		NO 177566 B	03-07-1995
		NO 941725 A,B,	27-09-1990
		NO 943057 A,B,	27-09-1990
		NO 951075 A,B,	27-09-1990
		NO 963245 A	27-09-1990
		NZ 228050 A	29-01-1992
		NZ 238843 A	29-01-1992
		NZ 238844 A	29-01-1992
		NZ 238845 A	29-01-1992
		PT 89774 A,B	04-10-1989
		US 5245047 A	14-09-1993
		US 5280126 A	18-01-1994
		WO 8907598 A	24-08-1989
		US 5124482 A	23-06-1992
		US 5149837 A	22-09-1992
		US 5216174 A	01-06-1993
		ZA 8900989 A	31-10-1990
		US 5097045 A	17-03-1992